

# ON THREEFOLDS WITHOUT NONCONSTANT REGULAR FUNCTIONS

JING ZHANG

**ABSTRACT.** We consider smooth threefolds  $Y$  defined over  $\mathbb{C}$  with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0, i > 0$ . Let  $X$  be a smooth projective threefold containing  $Y$  and  $D$  be the boundary divisor with support  $X - Y$ . We are interested in the following question: What geometry information of  $X$  can be obtained from the regular function information on  $Y$ ? Suppose that the boundary  $X - Y$  is a smooth projective surface. In this paper, we analyse two different cases, i.e., there are no nonconstant regular functions on  $Y$  or there are lots of regular functions on  $Y$ . More precisely, if  $H^0(Y, \mathcal{O}_Y) = \mathbb{C}$ , we prove that  $\frac{1}{2}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) \geq 0$ . In particular, if the line bundle  $\mathcal{O}_D(D)$  is not torsion, then  $q = h^1(X, \mathcal{O}_X) = 0$ ,  $\frac{1}{2}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) = 0$ ,  $\chi(\mathcal{O}_X) > 0$  and  $K_X$  is not nef. If there is a positive constant  $c$  such that  $h^0(X, \mathcal{O}_X(nD)) \geq cn^3$  for all sufficiently large  $n$  (we say that  $D$  is big or the  $D$ -dimension of  $X$  is 3) and  $D$  has no exceptional curves, then  $|nD|$  is base point free for  $n \gg 0$ . Therefore  $Y$  is affine if  $D$  is big.

## 1. INTRODUCTION

Let  $Y$  be an algebraic manifold (i.e., an irreducible smooth algebraic variety defined over  $\mathbb{C}$ ) with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ , where  $\Omega_Y^j$  is the sheaf of regular  $j$ -forms. We want to understand what  $Y$  is. This question was raised by J.-P. Serre for complex manifolds [Se]. Since  $Y$  is not compact, for any analytic or algebraic coherent sheaf  $\mathcal{F}$  on  $Y$ , we have  $H^3(Y, \mathcal{F}) = 0$  [Siu1, Siu2, Zh1]. So  $Y$  contains no complete surfaces [NS, Zh1]. If  $Y$  has non-constant regular functions, we know that it contains no complete curves [Zh1]. Let  $X$  be a smooth completion of  $Y$ , then the complement of  $Y$  in  $X$  is connected [Zh1]. Suppose that the boundary  $X - Y$  is of pure codimension 1 and is the support of an effective divisor  $D$  with simple normal crossings. We consider the  $D$ -dimension of  $X$  in order to understand  $Y$ . The notion of  $D$ -dimension is due to Iitaka ([I1], Lecture 3 or [Uen1], Chapter 2). It measures that how many regular functions there are on  $Y$ . If for all integers  $m > 0$  we have  $H^0(X, \mathcal{O}_X(mD)) = 0$ , then we define the  $D$ -dimension of  $X$ , denoted by  $\kappa(D, X)$ , to be  $-\infty$ . If  $h^0(X, \mathcal{O}_X(mD)) \geq 1$  for some  $m$ , choose a basis  $\{f_0, f_1, \dots, f_n\}$  of the linear space  $H^0(X, \mathcal{O}_X(mD))$ , it defines a rational map  $\Phi_{mD}$  from  $X$  to the projective space  $\mathbb{P}^n$  by sending a point  $x$  on  $X$  to  $(f_0(x), f_1(x), \dots, f_n(x))$  in  $\mathbb{P}^n$ . Then we define  $\kappa(D, X)$  to be the maximal dimension of the images of the rational map  $\Phi_{mD}$ , i.e.,

$$\kappa(D, X) = \max_m \{\dim(\Phi_{mD}(X))\}.$$

---

1991 *Mathematics Subject Classification.* Primary 14J30; Secondary 32Q28.

Let  $K_X$  be the canonical divisor of  $X$ , then the Kodaira dimension of  $X$  is the  $K_X$ -dimension of  $X$ , denoted by  $\kappa(X)$ , i.e.,

$$\kappa(X) = \kappa(K_X, X).$$

When  $\kappa(D, X) = \dim X$ , we say that  $D$  is big.

In our case, since  $D$  is effective, the  $D$ -dimension  $\kappa(D, X) \neq -\infty$ . In [Zh1, Zh2], we prove that  $\kappa(D, X)$  can be 1 and in this case, we have a surjective morphism from  $Y$  to a smooth affine curve  $C$  such that every fibre  $S$  satisfies the same vanishing condition, i.e.,  $H^i(S, \Omega_S^j) = 0$  for all  $j \geq 0$  and  $i > 0$ . We obtained that the Kodaira dimension of  $X$  is  $-\infty$  and  $q(X) = h^1(X, \mathcal{O}_X)$  can be any non-negative integer. In particular, all smooth fibres are of the same type, i.e., type (2) or type (3) open surface in the following Theorem 2.6 [Ku]. For the existence of the non-affine and non-product threefolds with vanishing Hodge cohomology, see [Zh2].

In [Zh3], we proved that  $\kappa(D, X) \neq 2$  and if  $\kappa(D, X) = 3$ , then  $Y$  is birational to  $\text{Spec} \Gamma(Y, \mathcal{O}_Y)$ . Furthermore, if  $Y$  is regularly separable, i.e., if for any two distinct points  $y_1$  and  $y_2$  on  $Y$ , there is a regular function  $f$  on  $Y$  such that  $f(y_1) \neq f(y_2)$ , then  $Y$  is affine. Regular separability implies  $\kappa(D, X) = 3$ . We want to know whether  $Y$  is affine if  $H^i(Y, \Omega_Y^j) = 0$  and  $\kappa(D, X) = 3$ . If we can prove the converse, that is,  $\kappa(D, X) = 3$  implies regular separability, then  $Y$  is affine. It is sufficient to prove that if  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$  and  $\kappa(D, X) = 3$ , then any non-constant regular function on  $Y$  defines an affine surface. Another possible approach is to prove that  $|nD|$  is base point free for some  $n > 0$ . Since  $Y$  contains no complete curves,  $Y$  is affine ([H2], Chapter 2, Proposition 2.2). In this paper, we will prove that when  $D$  is smooth, irreducible and contains no exceptional curves, then  $Y$  is affine.

**Theorem 1.1.** *Let  $Y$  be an irreducible smooth threefold contained in a smooth projective threefold  $X$  such that  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ . Suppose that the complement  $D = X - Y$  is a smooth projective surface without exceptional curves and  $\kappa(D, X) = 3$ , then  $|nD|$  is base point free for all  $n \gg 0$ .*

**Corollary 1.2.** *Under the condition of Theorem 1.1,  $Y$  is affine if and only if  $\kappa(D, X) = 3$ .*

The most mysterious case is  $\kappa(D, X) = 0$ . It is hard to understand because we cannot construct a fibre space by the divisor  $D$ . However, in order to keep track of  $Y$  and its cohomology, we have to use this boundary divisor to define the map. So we cannot apply Iitaka's fibration or Mori's construction. The situation on any normal and complete surface is much better. Any effective divisor on a normal complete surface has a unique Zariski decomposition and any two divisors have an intersection number [Sa2]. We even can get satisfied information of the surface if we know the numerical type of the divisor [Sa1]. When  $\dim Y = 3$  and the  $D$ -dimension is 0 or 3, we do not have Zariski decomposition [C] and a good method to check whether a divisor is nef. When  $\kappa(D, X) = 0$  and  $D$  is smooth and irreducible, we can reduce the problem to surface case.

**Theorem 1.3.** *Let  $Y$  be a smooth threefold contained in a smooth projective threefold  $X$  such that  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$  and the complement  $D = X - Y$  is a smooth projective surface. If  $Y$  has no nonconstant regular functions, then  $\frac{1}{2}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) \geq 0$ . In particular, if the line bundle  $\mathcal{O}_D(D)$  is not torsion, then  $q = h^1(X, \mathcal{O}_X) = 0$ ,  $\frac{1}{2}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) = 0$ ,  $\chi(\mathcal{O}_X) > 0$  and  $K_X$  is not nef.*

We organize the paper as follows. In Section 2, we will present some preparation. We will prove the two theorems in Section 3.

Let us mention some open problems for the Steinness of  $Y$ . When dimension of  $Y$  is 2, the type (3) open surfaces in Theorem 2.6 is a mystery. When dimension is 3 and  $\kappa(D, X) = 1$ , we do not know whether the nonaffine, nonproduct example in [Zh2] is Stein. Hartshorne asked the following question ([H2], page 235): Remove an irreducible curve from a smooth complete surface, what is the condition for the open surface to be Stein? See [N, Ued] for recent progress. Unfortunately, their approach cannot be applied to type (3) surface because the boundary has 9 components. We can ask three dimensional analogue of Hartshorne's question: What is the necessary and sufficient condition of a smooth threefold to be Stein but not affine? Are the two conditions  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$  and  $\kappa(D, X) \leq 1$  sufficient? The known result is that there exist algebraic Stein threefolds  $Y$  with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$ ,  $i > 0$  and  $\kappa(D, X) = 1$  [Zh1]. However, if  $\kappa(D, X) = 0$ , we do not know whether there exists an algebraic Stein threefold with  $H^i(Y, \Omega_Y^j) = 0$  whereas such surface exists ([H2], Chapter VI, Section 3 or [Ku]).

**Acknowledgments** I would like to thank the following professors for helpful discussions: Steven Dale Cutkosky, Dan Edidin, N.Mohan Kumar, Zhenbo Qin and Qi Zhang.

## 2. PREPARATION

**Lemma 2.1.** [Goodman, Hartshorne] *Let  $V$  be a scheme and  $D$  be an effective Cartier divisor on  $V$ . Let  $U = V - \text{Supp} D$  and  $F$  be any coherent sheaf on  $V$ , then for every  $i \geq 0$ ,*

$$\lim_{\vec{n}} H^i(V, F \otimes \mathcal{O}(nD)) \cong H^i(U, F|_U).$$

**Lemma 2.2.** *Let  $I = \{i\}$  be a direct system of indices. Let  $\{\mathcal{F}_i, f_{ji}\}$ ,  $\{\mathcal{G}_i, g_{ji}\}$  and  $\{\mathcal{H}_i, h_{ji}\}$  be direct system of coherent sheaves indexed by  $I$  over a topological space  $X$ . If for all  $i \in I$ , there are short exact sequences*

$$0 \longrightarrow \mathcal{F}_i \longrightarrow \mathcal{G}_i \longrightarrow \mathcal{H}_i \longrightarrow 0$$

*and the commutative diagram for all  $j \geq i$*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}_i & \longrightarrow & \mathcal{G}_i & \longrightarrow & \mathcal{H}_i \longrightarrow 0 \\ & & \downarrow f_{ji} & & \downarrow g_{ji} & & \downarrow h_{ji} \\ 0 & \longrightarrow & \mathcal{F}_j & \longrightarrow & \mathcal{G}_j & \longrightarrow & \mathcal{H}_j \longrightarrow 0, \end{array}$$

then we have exact sequence

$$0 \longrightarrow \lim_{\vec{i}} \mathcal{F}_i \longrightarrow \lim_{\vec{i}} \mathcal{G}_i \longrightarrow \lim_{\vec{i}} \mathcal{H}_i \longrightarrow 0.$$

*Proof.* By the assumption, for any point  $x \in X$ , we have short exact sequence on stalks

$$0 \longrightarrow (\mathcal{F}_i)_x \longrightarrow (\mathcal{G}_i)_x \longrightarrow (\mathcal{H}_i)_x \longrightarrow 0$$

and the commutative diagram for all  $j \geq i$

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\mathcal{F}_i)_x & \longrightarrow & (\mathcal{G}_i)_x & \longrightarrow & (\mathcal{H}_i)_x \longrightarrow 0 \\ & & \downarrow (f_{ji})_x & & \downarrow (g_{ji})_x & & \downarrow (h_{ji})_x \\ 0 & \longrightarrow & (\mathcal{F}_j)_x & \longrightarrow & (\mathcal{G}_j)_x & \longrightarrow & (\mathcal{H}_j)_x \longrightarrow 0. \end{array}$$

By Ueno, Algebraic Geometry 2 ([Uen2], Page 10), we have the following exact sequences of abelian groups on stalks

$$0 \longrightarrow \lim_{\vec{i}} (\mathcal{F}_i)_x \longrightarrow \lim_{\vec{i}} (\mathcal{G}_i)_x \longrightarrow \lim_{\vec{i}} (\mathcal{H}_i)_x \longrightarrow 0.$$

Since direct limits commute with each other ([Br], page 20), we have

$$\lim_{\vec{i}} (\mathcal{F}_i)_x = \lim_{\vec{i}} \lim_{x \in U} \mathcal{F}_i = \lim_{x \in U} \lim_{\vec{i}} \mathcal{F}_i = (\lim_{\vec{i}} \mathcal{F}_i)_x.$$

The Lemma follows from

$$0 \longrightarrow (\lim_{\vec{i}} \mathcal{F}_i)_x \longrightarrow (\lim_{\vec{i}} \mathcal{G}_i)_x \longrightarrow (\lim_{\vec{i}} \mathcal{H}_i)_x \longrightarrow 0.$$

Q.E.D.

**Lemma 2.3.** *Let  $Y$  be a smooth variety contained in a smooth projective variety  $X$  such that the complement  $X - Y$  is compact and of pure codimension 1. Let  $D$  be any effective divisor with support  $X - Y$ . Then we have the following two exact sequences*

$$0 \longrightarrow \lim_{\vec{n}} \mathcal{O}_X(nD) \longrightarrow \lim_{\vec{n}} \mathcal{O}_X((n+1)D) \longrightarrow \lim_{\vec{n}} \mathcal{O}_D((n+1)D) \longrightarrow 0$$

and for all  $j > 0$

$$0 \longrightarrow \lim_{\vec{n}} \Omega_X^j(nD) \longrightarrow \lim_{\vec{n}} \Omega_X^j((n+1)D) \longrightarrow \lim_{\vec{n}} \Omega_X^j((n+1)D)|_D \longrightarrow 0.$$

*Proof.* For any positive integers  $n$  and  $m$ , we have the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X(nD) & \xrightarrow{f} & \mathcal{O}_X((n+1)D) & \xrightarrow{r} & \mathcal{O}_D((n+1)D) \longrightarrow 0 \\ & & \downarrow i & & \downarrow i & & \downarrow h \\ 0 & \longrightarrow & \mathcal{O}_X((n+m)D) & \xrightarrow{f} & \mathcal{O}_X((n+m+1)D) & \xrightarrow{r} & \mathcal{O}_D((n+m+1)D) \longrightarrow 0, \end{array}$$

where the first map  $f$  in each arrow is defined by the local defining function of  $D$ ,  $r$  is the restriction map, the vertical map  $i$  is the natural embedding map "1" and the last vertical map  $h$  is defined as follows. For any nonzero element  $s$  in  $\mathcal{O}_D((n+1)D)$  (locally), there is an  $t$  in  $\mathcal{O}_X((n+1)D)$  (locally) such that  $r(t) = s$ .

Then we define  $h(s)$  to be  $r(i(t)) = r(t)$ . Since  $i \neq f$ , if  $t$  is not zero on  $D$ , then  $i(t)$  does not sit in the image of  $\mathcal{O}_X((n+m)D)$  under the map  $f$ . Notice that the vertical map  $i$  defines the direct limit. Then the first exact sequence is an immediate consequence of Lemma 2.2. The second exact sequence can be similarly proved.

Q.E.D.

**Lemma 2.4.** *Let  $Y$  be a smooth threefold with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ . Let  $X$  be a smooth completion of  $Y$  such that the complement  $X - Y$  is compact and of pure codimension 1. Let  $D$  be any effective divisor on  $X$  with support  $X - Y$ , then for all  $j \geq 0$  and  $i > 0$ , we have*

$$\lim_{\vec{n}} H^i(D, \Omega_X^j(nD)|_D) = 0.$$

*Proof.* Since  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ , by Lemma 2.1, we have

$$\lim_{\vec{n}} H^i(X, \Omega_X^j(nD)) = 0.$$

Since the direct limit commutes with cohomology ([H1], Chapter III, Proposition 2.9), the Lemma follows from the short exact sequences in Lemma 2.3.

Q.E.D.

The following lemma is not new. We already proved it in our previous paper [Zh1]. Since we frequently use the argument in the proof of the theorems, We include a proof here for completeness.

**Lemma 2.5.** *Under the condition of Lemma 2.4,  $H^3(X, \mathcal{O}_X(nD)) = 0$  for sufficiently large  $n$ .*

*Proof.* From the exact sequence

$$0 \longrightarrow \mathcal{O}_X(nD) \longrightarrow \mathcal{O}_X((n+1)D) \longrightarrow \mathcal{O}_D((n+1)D) \longrightarrow 0,$$

we have surjective map from  $H^3(X, \mathcal{O}_X(nD))$  to  $H^3(X, \mathcal{O}_X((n+1)D))$  for all  $n \geq 0$ . By Lemma 2.1, we have

$$\lim_{\vec{n}} H^3(X, \mathcal{O}_X(nD)) = 0.$$

These two conditions imply the vanishing of the third cohomology for large  $n$ .

Q.E.D.

**Theorem 2.6.** [Mohan Kumar] *Let  $Y$  be a smooth algebraic surface over  $\mathbb{C}$  with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ , then  $Y$  is one of the following*

- (1)  $Y$  is affine.
- (2) Let  $C$  be an elliptic curve and  $E$  the unique nonsplit extension of  $\mathcal{O}_C$  by itself. Let  $X = \mathbb{P}_C(E)$  and  $D$  be the canonical section, then  $Y = X - D$ .
- (3) Let  $X$  be a projective rational surface with an effective divisor  $D = -K$  with  $D^2 = 0$ ,  $\mathcal{O}(D)|_D$  be nontorsion and the dual graph of  $D$  be  $\tilde{D}_8$  or  $\tilde{E}_8$ , then  $Y = X - D$ .

In the above theorem, when  $Y$  is affine, we can choose  $D$  such that  $D$  is ample ([H2], Theorem 4.2, page 69), so  $\kappa(D, X) = 2$ . If  $Y$  is not affine, then  $\kappa(D, X) = 0$  by Lemma 1.8 [Ku].

**Theorem 2.7. [Cutkosky, Srinivas]** *Let  $k$  be a field of characteristic 0. Let  $X$  be a normal surface, proper over  $k$ , and let  $D$  be an effective Cartier divisor on  $X$ . Then, for sufficiently large  $n$ ,*

$$h^0(X, \mathcal{O}_X(nD)) = P(n) + \lambda(n),$$

where  $P(n)$  is a quadratic polynomial and the function  $\lambda(n)$  is periodic.

For the proof of the following Iitaka's theorem, see Lecture 3 [I1] or Theorem 8.1 [Uen1].

**Theorem 2.8. [Iitaka]** *Let  $X$  be a normal projective variety and let  $D$  be an effective divisor on  $X$ . There exist two positive numbers  $\alpha$  and  $\beta$  such that for all sufficiently large  $n$  we have*

$$\alpha n^{\kappa(D, X)} \leq h^0(X, \mathcal{O}_X(nD)) \leq \beta n^{\kappa(D, X)}.$$

### 3. PROOF OF THE THEOREMS

From now on, we will fix the notations as follows.  $Y$  is a smooth threefold with  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$ .  $X$  is a smooth completion of  $Y$  such that the complement  $X - Y = D$  is a smooth projective surface.

**Proof of Theorem 1.1.** Since  $\kappa(D, X) = 3$  and  $D$  is effective, by Theorem 2.8, there is a constant  $c > 0$  such that for sufficiently large  $n$ ,  $h^0(X, \mathcal{O}_X(nD)) > cn^3$ . From the short exact sequence

$$0 \longrightarrow \mathcal{O}_X(nD) \longrightarrow \mathcal{O}_X((n+1)D) \longrightarrow \mathcal{O}_D((n+1)D) \longrightarrow 0,$$

we have

$$0 \longrightarrow H^0(\mathcal{O}_X(nD)) \longrightarrow H^0(\mathcal{O}_X((n+1)D)) \longrightarrow H^0(\mathcal{O}_D((n+1)D)) \longrightarrow \dots$$

So there are infinitely many  $n$ , such that  $h^0(\mathcal{O}_D(nD)) > 0$  since  $h^0(X, \mathcal{O}_X(nD))$  grows like  $cn^3$  for sufficiently large  $n$ . Let

$$N = \{m \in \mathbb{N}, h^0(D, \mathcal{O}_D(mD)) > 0\}.$$

Let  $p$  be the greatest common divisor of  $N$ , then

$$h^0(D, \mathcal{O}_D(npD)) > 0$$

for all  $n \geq 0$ . Thus the line bundle  $\mathcal{O}_D(pD)$  determines an effective divisor  $G$  on  $D$  such that

$$\mathcal{O}_D(pD) = \mathcal{O}_D(G).$$

If  $\mathcal{O}_D(D)$  is torsion, then by Lecture 3, [I1],  $h^0(D, \mathcal{O}_D(nD)) \leq 1$  for all  $n \geq 0$ . Since  $D$  is big, i.e.,  $h^0(X, \mathcal{O}_X(nD)) > cn^3$ ,  $\mathcal{O}_D(D)$  is not torsion. So we may assume that  $D|_D$  determines an effective divisor on  $D$ . We still denote it as  $G$ . By Lemma 2.4, we have for all  $j \geq 0$  and  $i > 0$ ,

$$\lim_{\vec{n}} H^i(D, \Omega_X^j(nD)|_D) = 0.$$

In particular, for all  $j \geq 0$  and  $i > 0$ ,

$$\begin{aligned} \lim_{\vec{n}} H^i(D, \Omega_X^j(nD)|_D) &= \lim_{\vec{n}} H^i(D, \Omega_X^j \otimes \mathcal{O}_D(nD)) \\ &= \lim_{\vec{n}} H^i(D, \Omega_X^j \otimes \mathcal{O}_D(nG)) = 0. \end{aligned}$$

Let  $S = D - G$ , by Lemma 2.1, for all  $i > 0$  and  $j \geq 0$ , we have

$$H^i(S, \Omega_X^j|_S) = H^i(S, \Omega_D^j|_S) = 0.$$

By the first exact sequence of Lemma 2.3, we have

$$\begin{aligned} \cdots \longrightarrow H^1(X, \lim_{\vec{n}} \mathcal{O}_X((n+1)D)) &\longrightarrow H^1(D, \lim_{\vec{n}} \mathcal{O}_D((n+1)D)) \\ &\longrightarrow H^2(X, \lim_{\vec{n}} \mathcal{O}_X(nD)) \longrightarrow \cdots \end{aligned}$$

Since the direct limit commutes with cohomology [H1], Chapter III, Proposition 2.9, and by Lemma 2.1, for all  $i > 0$ ,

$$\lim_{\vec{n}} H^i(X, \mathcal{O}_X(nD)) = 0,$$

we have

$$H^i(S, \mathcal{O}_S) = \lim_{\vec{n}} H^i(D, \mathcal{O}_D(nD)) = 0.$$

By [H1], page 178, Theorem 8.17 or [GrH], page 157, (\*\*), we have

$$0 \longrightarrow \mathcal{O}_D(-D) \longrightarrow \Omega_X^1|_D \longrightarrow \Omega_D^1 \longrightarrow 0.$$

Tensoring with  $\mathcal{O}_D((n+1)D)$ , we have

$$0 \longrightarrow \mathcal{O}_D(nD) \longrightarrow \Omega_X^1|_D \otimes \mathcal{O}_D((n+1)D) \longrightarrow \Omega_D^1((n+1)D) \longrightarrow 0.$$

Taking the direct limit, the corresponding long exact sequence gives

$$H^i(S, \Omega_S^1) = \lim_{\vec{n}} H^i(D, \Omega_D^1((n+1)D)) = 0.$$

Applying the same procedure to the following short exact sequence

$$0 \longrightarrow \Omega_D^1(-D) \longrightarrow \Omega_X^2|_D \longrightarrow \Omega_D^2 \longrightarrow 0,$$

we have  $H^i(S, \Omega_S^2) = 0$ . So  $S$  satisfies the same vanishing condition, i.e., for all  $i > 0$ ,  $j \geq 0$ ,  $H^i(S, \Omega_S^j) = 0$ .

By Theorem 2.6 in Section 2 and Lemma 1.8 [Ku], the boundary divisor  $G$  on  $D$  is connected and  $\kappa(G, D) = 0$  or  $2$ . We know that there are  $(-2)$ -curves on type (3) surface in Theorem 2.6. Since  $D$  has no exceptional curves,  $D$  is not of type (3). Let  $C$  be an irreducible complete curve on  $X$ . If  $C$  is not contained in  $D$ , then  $C \cdot D > 0$  since  $Y$  has no complete curves by Lemma 5 [Zh1]. If  $C$  is contained in  $D$  but not in  $G$ , then again

$$C \cdot D = C \cdot D|_D = C \cdot G > 0$$

since  $S$  has no complete curves by Lemma 1.1 [Ku]. If  $C$  is a component of  $G$ , then  $C \cdot G \geq 0$  since  $G$  has no exceptional curves and is connected. Therefore  $D$  is nef. We know that  $S$  is either affine or type (2) surface in Theorem 2.6. If  $S$  is of type

(2), then the boundary  $D - S$  is an irreducible elliptic curve with self-intersection number 0 by Theorem 2.6. This gives us  $G^2 = D^3 = 0$  which is impossible since  $D$  is nef and  $h^0(X, \mathcal{O}_X(nD)) > cn^3$  (Proposition 2.61, [KM]). So  $S$  must be affine. Therefore for any component  $C$  of  $G$ ,

$$D^3 = G^2 \geq C \cdot G > 0.$$

Therefore  $D$  is big and nef.

We will prove that the linear system  $|nD|$  is base point free for sufficiently large  $n$ .

Since  $S$  is an affine surface and  $G$  contains no exceptional curves,  $G$  is ample, i.e.,  $pD|_D$  is ample on  $D$ . Thus  $D|_D$  is ample on  $D$ . Let  $L = D|_D$ , then

$$H^1(D, \mathcal{O}_D(nL)) = H^1(D, \mathcal{O}_D(nD)) = 0$$

for sufficiently large  $n$ . Thus for all sufficiently large  $n$  we have surjective map

$$H^1(X, \mathcal{O}_X((n-1)D)) \longrightarrow H^1(X, \mathcal{O}_X(nD)) \longrightarrow 0.$$

Since

$$\lim_{\vec{n}} H^1(X, \mathcal{O}_X(nD)) = 0,$$

we have  $H^1(X, \mathcal{O}_X(nD)) = 0$  for large  $n$ . Thus we have an exact sequence

$$0 \longrightarrow H^0(X, \mathcal{O}_X((n-1)D)) \longrightarrow H^0(X, \mathcal{O}_X(nD)) \longrightarrow H^0(D, \mathcal{O}_D(nD)) \longrightarrow 0.$$

This implies that the linear system  $|nD|$  is base point free. In fact, a point  $x \in X$  is a base point of  $|nD|$  if and only if for every section  $s \in H^0(\mathcal{O}_X(nD))$ ,  $s(x) = 0$  or if and only if for every effective divisor  $E \in |nD|$ ,  $x \in E$ . Suppose that  $x$  is a base point of  $|nD|$ , then  $x \in nD$ . Thus  $x \in D$ . Since  $nD|_D$  is very ample for large  $n$ , there is a divisor  $F \in |nD|_D$  such that  $x$  is not a point of  $F$ . Pull  $F$  back to  $|nD|$ , then there is a divisor  $E \in |nD|$ , such that  $x$  is not a point of  $E$ . This is a contradiction. So the linear system  $|nD|$  is base point free for sufficiently large  $n$ .

Q.E.D.

**Proof of Theorem 1.3.** In order to prove Theorem 1.3, we will analyse three cases:  $D^3 > 0$ ,  $D^3 < 0$  and  $D^3 = 0$ .

**Proposition 3.1.** *Let  $Y$  be a smooth threefold contained in a smooth projective threefold  $X$  such that  $H^0(Y, \mathcal{O}_Y) = \mathbb{C}$ ,  $H^i(Y, \Omega_Y^j) = 0$  for all  $j \geq 0$  and  $i > 0$  and the complement  $D = X - Y$  is a smooth projective surface. Then  $D^3$  is not positive.*

*Proof.* Suppose  $D^3 > 0$ . By the Riemann-Roch formula for surfaces, we have

$$h^0(\mathcal{O}_D(nD)) - h^1(\mathcal{O}_D(nD)) + h^2(\mathcal{O}_D(nD)) = \chi(\mathcal{O}_D) + \frac{1}{2}n^2D^3 - \frac{1}{2}nD^2K_D,$$

$$h^0(\mathcal{O}_D(-nD)) - h^1(\mathcal{O}_D(-nD)) + h^2(\mathcal{O}_D(-nD)) = \chi(\mathcal{O}_D) + \frac{1}{2}n^2D^3 + \frac{1}{2}nD^2K_D.$$

So we have

$$h^0(\mathcal{O}_D(nD)) + h^2(\mathcal{O}_D(nD)) \geq \chi(\mathcal{O}_D) + \frac{1}{2}n^2D^3 - \frac{1}{2}nD^2K_D,$$



$$h^0(\mathcal{O}_D(-nD)) + h^2(\mathcal{O}_D(-nD)) \geq \chi(\mathcal{O}_D) + \frac{1}{2}n^2D^3 + \frac{1}{2}nD^2K_D.$$

Since  $D^3 > 0$  and  $(K_D + nD|_D) + (K_D - nD|_D) = 2K_D$ , either  $h^0(\mathcal{O}_D(nD)) \rightarrow \infty$  or  $h^0(\mathcal{O}_D(-nD)) \rightarrow \infty$  as  $n \rightarrow \infty$  and they cannot be big simultaneously. Otherwise, we would have  $h^0(\mathcal{O}_D(2K_D)) = \infty$  which is absurd.

Case 1.  $h^0(\mathcal{O}_D(nD)) \rightarrow \infty$ .

In this case, there is an integer  $n_0 > 0$  such that for all  $n \geq n_0$ ,  $nD|_D$  is an effective divisor on  $D$ . We may assume that  $D|_D = G$  is effective without loss of generality. From the exact sequence

$$0 \rightarrow \mathcal{O}_D(nG) \rightarrow \mathcal{O}_D((n+1)G) \rightarrow \mathcal{O}_G((n+1)G) \rightarrow 0,$$

since  $H^2(\mathcal{O}_G((n+1)G)) = 0$ , we have surjective map

$$H^2(\mathcal{O}_D(nG)) \rightarrow H^2(\mathcal{O}_D((n+1)G)) \rightarrow 0.$$

By Lemma 2.4,

$$\lim_{\vec{n}} H^2(\mathcal{O}_D(nG)) = \lim_{\vec{n}} H^2(\mathcal{O}_D(nD)) = 0.$$

So for all sufficiently large  $n$ ,  $H^2(\mathcal{O}_D(nD)) = 0$ . By the same argument,  $H^2(\mathcal{O}_X(nD)) = 0$  for  $n \gg 0$ . Since by Lemma 2.5,  $H^3(\mathcal{O}_X(nD)) = 0$ , we have

$$\chi(\mathcal{O}_X(nD)) = h^0(\mathcal{O}_X(nD)) - h^1(\mathcal{O}_X(nD)) = 1 - h^1(\mathcal{O}_X(nD)) \leq 1.$$

On the other hand, since  $D^3 > 0$ , by the Riemann-Rock formula for threefolds ([Ful], page 291), we have

$$\chi(\mathcal{O}_X(nD)) = \frac{1}{6}(nD)^3 + \frac{1}{4}c_1 \cdot (nD)^2 + \frac{1}{12}(c_1^2 + c_2) \cdot nD + \frac{1}{24}c_1 \cdot c_2 \rightarrow \infty$$

as  $n \rightarrow \infty$ . Therefore we get a contradiction. Thus  $h^0(\mathcal{O}_D(nD)) \rightarrow \infty$  is not a possible case.

Case 2.  $h^0(\mathcal{O}_D(-nD)) \rightarrow \infty$  as  $n \rightarrow \infty$ .

There is an  $n_0 > 0$  such that for all  $n \geq n_0$ ,  $-nD|_D$  is effective divisor on  $D$ . Again we may assume that  $G = -D|_D$  is effective. Since  $(K_X + D)|_D = K_D$  (Proposition 8.20, Chapter II, [H1]), by Lemma 2.4 and Serre duality, we have

$$\begin{aligned} \lim_{\vec{n}} H^2(D, \Omega_X^3((n+1)D)|_D) &= \lim_{\vec{n}} H^2(D, \mathcal{O}_X(K_X + (n+1)D)|_D) \\ &= \lim_{\vec{n}} H^2(D, \mathcal{O}_D(K_D + nD|_D)) = 0. \end{aligned}$$

Take the duality of the vector space, we have ([H2], Chapter 3, Section 3)

$$\lim_{\vec{n}} H^0(D, \mathcal{O}_D(-nD)) = (\lim_{\vec{n}} H^2(D, \mathcal{O}_D(K_D + nD|_D)))^* = 0,$$

where  $*$  indicates the dual vector space. Let  $A_n = H^0(\mathcal{O}_D)$ , then  $A_n = \mathbb{C}$ . Let  $B_n = H^0(D, \mathcal{O}_D(-nD))$  and let  $C_n$  be their quotient, then we have short exact sequence

$$0 \rightarrow A_n \rightarrow B_n \rightarrow C_n \rightarrow 0.$$

Since  $-D|_D = G$  is effective,  $B_n$  is a subspace of linear space  $B_{n'}$  if  $n' > n$ . The map  $B_{n'} \rightarrow B_n$  is the natural restriction map. So we have the exact sequence of inverse systems

$$0 \longrightarrow (A_n) \longrightarrow (B_n) \longrightarrow (C_n) \longrightarrow 0.$$

By [H1], page 191, we have injective map

$$0 \longrightarrow \mathbb{C} = \varprojlim A_n \longrightarrow \varprojlim B_n$$

which is contrary to

$$\varprojlim B_n = 0.$$

Thus  $h^0(\mathcal{O}_D(-nD)) \longrightarrow \infty$  as  $n \longrightarrow \infty$  is also impossible.

We have seen that  $D^3$  is not positive.

Q.E.D.

*Remark 3.2.* Since  $h^0(X, \mathcal{O}_X(nD)) = 1$ , when  $D$  is not irreducible but nef, we still have  $D^3 \leq 0$  [KM], Proposition 2.61.

**Proposition 3.3.** *With the assumption of Proposition 3.1,  $D^3$  is not negative.*

*Proof.* Suppose  $D^3 < 0$ . If  $h^0(\mathcal{O}_D(nD)) = 0$  for all  $n \gg 0$ , then we have injective map

$$0 \longrightarrow H^1(X, \mathcal{O}_X(nD)) \longrightarrow H^1(X, \mathcal{O}_X((n+1)D)).$$

By Lemma 2.4,

$$\varprojlim H^1(X, \mathcal{O}_X(nD)) = 0.$$

These two restrictions imply  $H^1(X, \mathcal{O}_X(nD)) = 0$  for all  $n \gg 0$ . Since  $H^3(X, \mathcal{O}_X(nD)) = 0$  for all  $n \gg 0$ , the Euler characteristic

$$\chi(\mathcal{O}_X(nD)) = 1 + h^2(\mathcal{O}_X(nD)) \geq 1.$$

On the other hand, since  $D^3 < 0$ , by the Riemann-Roch formula ([Ful], page 291), we have

$$\begin{aligned} \chi(\mathcal{O}_X(nD)) &= \frac{1}{6}n^3D^3 + \frac{1}{4}c_1 \cdot (nD)^2 + \frac{1}{12}(c_1^2 + c_2) \cdot nD + \frac{1}{24}c_1 \cdot c_2 \\ &\longrightarrow -\infty \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This contradicts the fact that  $\chi(\mathcal{O}_D(nD))$  is positive. Therefore there are infinitely many  $n$  such that  $h^0(\mathcal{O}_D(nD)) > 0$ . Let  $N, p$  be the same as in the proof of Theorem 1.1, i.e.,

$$N = \{m \in \mathbb{N}, h^0(D, \mathcal{O}_D(mD)) > 0\},$$

and  $p$  is the greatest common divisor of  $N$ . Then  $h^0(D, \mathcal{O}_D(pD)) > 0$ . Since  $D^3 \neq 0$ , the line bundle  $\mathcal{O}_D(pD)$  is not trivial and  $pD|_D$  defines an effective divisor on surface  $D$ . Thus we may assume that  $D|_D = G$  is an effective divisor on  $D$ .

Let  $S = D - G$ , by the same argument as in the proof of Theorem 1.1,  $H^i(S, \Omega_S^j) = 0$  for all  $i > 0$  and  $j \geq 0$  since

$$\varprojlim H^i(D, \Omega_X^j(nD)|_D) = 0.$$

By Theorem 2.6 in Section 2 and Lemma 1.8 [Ku],  $\kappa(G, D) = 0$  or  $2$ . If  $\kappa(G, D) = 2$ , then by Theorem 2.7, for all  $n \gg 0$ ,

$$h^0(D, \mathcal{O}_D(nG)) = a_2 n^2 + a_1 n + a_0 + \lambda(n),$$

where  $\lambda(n)$  is periodic. By Theorem 2.8, since  $\kappa(G, D) = 2$ , there is a constant  $c > 0$  such that  $h^0(D, \mathcal{O}_D(nG)) > cn^2$ . Thus  $a_2 > 0$ . By Lemma 2.1,

$$\lim_{\vec{n}} H^0(D, \mathcal{O}_D(nG)) = \lim_{\vec{n}} H^0(D, \mathcal{O}_D(nD)) = H^0(S, \mathcal{O}_S) \neq 0.$$

In fact, since  $S$  is affine,  $h^0(S, \mathcal{O}_S) = \infty$ . By Lemma 2.3, we have

$$\begin{aligned} 0 \longrightarrow \lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD - D)) &\longrightarrow \lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD)) \\ &\longrightarrow \lim_{\vec{n}} H^0(X, \mathcal{O}_D(nD)) \longrightarrow 0. \end{aligned}$$

By Lemma 2.1,

$$\lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD)) = H^0(Y, \mathcal{O}_Y) = \mathbb{C}$$

and

$$\lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD - D)) = H^0(Y, \mathcal{O}_X(-D)|_Y) = \mathbb{C}.$$

Thus the above exact sequence is

$$0 \longrightarrow \mathbb{C} \longrightarrow \mathbb{C} \longrightarrow H^0(S, \mathcal{O}_S) \longrightarrow 0.$$

This is impossible since  $h^0(S, \mathcal{O}_S) = \infty$ . So  $\kappa(G, D) \neq 2$ .

If  $\kappa(G, D) = 0$ , since  $\mathcal{O}_D(D) \neq \mathcal{O}_D$  defines an effective divisor on  $D$ , by Lemma 2.1, 2.4, and Lemma 1.8[Ku] we have

$$\lim_{\vec{n}} H^0(D, \mathcal{O}_D(nD)) = H^0(S, \mathcal{O}_S) = \mathbb{C}.$$

Again by Lemma 2.1 and 2.3, we have

$$\begin{aligned} 0 \longrightarrow \lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD - D)) &\longrightarrow \lim_{\vec{n}} H^0(X, \mathcal{O}_X(nD)) \\ &\longrightarrow \lim_{\vec{n}} H^0(D, \mathcal{O}_D(nD)) \longrightarrow 0, \end{aligned}$$

which is impossible since three direct limits are  $\mathbb{C}$ . Therefore  $D^3$  is not negative.

Q.E.D.

**Proposition 3.4.** *Under the condition of Proposition 3.1,  $D^3 = 0$  and  $D|_D$  is numerically equivalent to 0.*

*Proof.* Let  $H$  be an ample divisor on  $X$ . Then when restricted on  $D$ ,  $L = H|_D$  is still ample by Proposition 4.1 [H2]. We may assume that  $L$  is smooth by Bertini's Theorem (Chapter 2, Theorem 8.8 [H1] or Theorem 4.21, [Uen1]). Let  $G$  be the divisor determined by the line bundle  $\mathcal{O}_D(D)$ .

If  $L \cdot G \neq 0$ , then there exists an  $n_0 > 0$  such that either the linear system  $|n_0 G|$  is nonempty or  $|-n_0 G|$  is nonempty. If  $|n_0 G|$  is nonempty, we may assume that  $G$  is effective. Let  $S = D - G$ . By Lemma 2.3, we have

$$\lim_{\vec{n}} H^0(X, \mathcal{O}_D(nD)) = H^0(S, \mathcal{O}_S).$$

By Theorem 2.8,

$$h^0(S, \mathcal{O}_S) = \dim_{\mathbb{C}} \lim_{\vec{n}} H^0(X, \mathcal{O}_D(nD)) \neq 0.$$

By the first exact sequence in Lemma 2.3, we have exact sequence

$$0 \longrightarrow \mathbb{C} \longrightarrow \mathbb{C} \longrightarrow H^0(S, \mathcal{O}_S) \longrightarrow 0.$$

This is impossible.

If  $|-n_0 G|$  is nonempty, then we may assume that  $-G$  is an effective divisor on  $D$ . By the same argument as in case 2, Proposition 3.1, we have

$$\lim_{\vec{n}} H^0(D, \mathcal{O}_D(-nD)) = (\lim_{\vec{n}} H^2(D, \mathcal{O}_D(K_D + nD)))^* = 0,$$

which is again not possible since

$$\mathbb{C} = \lim_{\vec{n}} H^0(D, \mathcal{O}_D) \hookrightarrow \lim_{\vec{n}} H^0(D, \mathcal{O}_D(-nD)).$$

We have seen that the only possible case is  $L \cdot G = 0$ . Since  $G^2 = D^3 = 0$ , by the Hodge Index Theorem,  $G$  is numerically equivalent to 0. If  $G = 0$ , then

$$(\lim_{\vec{n}} H^2(D, \mathcal{O}_D(K_D + nD)))^* = \lim_{\vec{n}} H^0(D, \mathcal{O}_D(-nD)) = \mathbb{C}$$

which contradicts Lemma 2.4. So we have  $G \equiv 0$  but  $G \neq 0$ .

Q.E.D.

**Proposition 3.5.** *Under the condition of Proposition 3.1, If  $\mathcal{O}_D(D)$  is not torsion, then  $q = h^1(X, \mathcal{O}_X) = 0$ ,  $\frac{1}{2}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) = 0$ ,  $\chi(\mathcal{O}_X) > 0$  and  $K_X$  is not nef.*

*Proof.* If there is an  $n_0$  such that  $h^0(D, \mathcal{O}_D(n_0 D)) \neq 0$  (or  $h^0(D, \mathcal{O}_D(-n_0 D)) \neq 0$ ), then  $\kappa(D, D|_D) \geq 0$ . By [I1], page 34-35, there are infinitely many  $n$  such that  $h^0(D, \mathcal{O}_D(nD)) \neq 0$  (or  $h^0(D, \mathcal{O}_D(-nD)) \neq 0$ ). Then  $D|_D$  (or  $-D|_D$ ) defines an effective divisor on  $D$  since  $\mathcal{O}_D(D)$  is not torsion. By Lemma 2.3, we can get a contradiction

$$0 \longrightarrow \mathbb{C} \longrightarrow \mathbb{C} \longrightarrow H^0(S, \mathcal{O}_S) \longrightarrow 0.$$

So  $h^0(D, \mathcal{O}_D(nD)) = h^0(D, \mathcal{O}_D(-nD)) = 0$  for all  $n > 0$ .

Since  $h^0(D, \mathcal{O}_D(nD)) = 0$  for all  $n > 0$ , from the exact sequence

$$0 \longrightarrow \mathcal{O}_X((n-1)D) \longrightarrow \mathcal{O}_X(nD) \longrightarrow \mathcal{O}_D(nD) \longrightarrow 0,$$

we have injective map from  $H^1(X, \mathcal{O}_X(nD))$  to  $H^1(X, \mathcal{O}_X((n+1)D))$ . Since the direct limit is 0,  $H^1(X, \mathcal{O}_X(nD)) = 0$  for sufficiently large  $n$ . By the injectivity,

$H^1(X, \mathcal{O}_X((n-1)D)) = 0$  for all  $n > 0$ , i.e.,  $H^1(X, \mathcal{O}_X(nD)) = H^1(X, \mathcal{O}_X) = 0$ . Thus for all  $n > 0$ , by Lemma 2.5, we have exact sequence

$$0 \longrightarrow H^1(D, \mathcal{O}_D((n+1)D)) \longrightarrow H^2(X, \mathcal{O}_X(nD)) \longrightarrow H^2(X, \mathcal{O}_X((n+1)D)) \\ \longrightarrow H^2(D, \mathcal{O}_D((n+1)D)) \longrightarrow 0.$$

This gives us

$$h^2(\mathcal{O}_X((n+1)D)) - h^2(\mathcal{O}_X(nD)) = h^2(\mathcal{O}_D((n+1)D)) - h^1(\mathcal{O}_D((n+1)D)).$$

By Riemann-Roch formulas for surfaces and threefolds, we have

$$\frac{1}{12}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D).$$

Since for all  $n \gg 0$ ,  $h^1(\mathcal{O}_X(nD)) = h^3(\mathcal{O}_X(nD)) = 0$  and  $D^3 = D^2 \cdot K_D = 0$ , we have  $\chi(\mathcal{O}_X(nD)) = 1 + h^2(\mathcal{O}_X(nD)) \geq 1$ . So  $\frac{1}{12}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) \geq 0$ .

If  $n \gg 0$ ,  $h^1(\mathcal{O}_D(nD)) = 0$ , then by the above exact sequence, we have

$$0 \longrightarrow H^2(X, \mathcal{O}_X(nD)) \longrightarrow H^2(X, \mathcal{O}_X((n+1)D)) \longrightarrow H^2(D, \mathcal{O}_D((n+1)D)) \longrightarrow 0,$$

It is easy to see that for all  $n \geq 0$ ,  $h^2(X, \mathcal{O}_X(nD)) = 0$  and  $h^2(\mathcal{O}_D((n+1)D)) = h^1(\mathcal{O}_D((n+1)D)) = 0$ . Thus  $\chi(\mathcal{O}_D) = 0$  and the proposition follows.

So the remaining case is: for infinitely many  $n > 0$ ,  $h^1(\mathcal{O}_D(nD)) > 0$ . We claim that  $\chi(\mathcal{O}_D) = 0$ . If not, then  $\chi(\mathcal{O}_D) > 0$ . Since for all  $n > 0$ ,  $h^0(D, \mathcal{O}_D(-nD)) = h^2(D, \mathcal{O}_D(K_D + nD)) = 0$ , we have surjective map from  $H^2(X, \mathcal{O}_X(K_X + (n-1)D))$  to  $H^2(X, \mathcal{O}_X(K_X + nD))$ . By Lemma 2.4,

$$\lim_{\vec{n}} H^2(X, \mathcal{O}_X(K_X + nD)) = 0.$$

So for  $n \gg 0$ ,  $H^2(X, \mathcal{O}_X(K_X + nD)) = 0$ .

Applying Riemann-Roch to the sheaf  $\mathcal{O}_X(K_X + nD)$ , we have

$$h^0(X, \mathcal{O}_X(K_X + nD)) \geq (\frac{1}{12}(c_1^2 + c_2) \cdot D)n > 0.$$

Since  $h^0(X, \mathcal{O}_X(nD)) = 1$  for all  $n \geq 0$ , the above inequality is not true. To see this, let  $D_m$  be an effective divisor in the linear system  $|K_X + mD|$  and  $D_{m+n}$  be an effective divisor in the linear system  $|K_X + (m+n)D|$ , then  $D_m + nD$  is linearly equivalent to  $D_{m+n}$ . Let  $f$  be the nonconstant rational function on  $X$  such that  $\text{div} f + D_{m+n} = D_m + nD$ , then  $\text{div} f + D_{m+n} - D_m = nD > 0$ . This gives  $f \in H^0(X, \mathcal{O}_X(D_{m+n} - D_m))$  and  $D_{m+n} - D_m$  is linearly equivalent to  $nD$ . Since  $h^0(X, \mathcal{O}_X(K_X + (m+n)D)) \geq c(m+n)$ , there are at least 2 linearly independent  $f$  in  $H^0(X, \mathcal{O}_X(D_{m+n} - D_m))$ . It contradicts to  $H^0(X, \mathcal{O}_X(D_{m+n} - D_m)) = H^0(X, \mathcal{O}_X(nD)) = \mathbb{C}$  for all  $n \geq 0$ . Thus  $h^0(X, \mathcal{O}_X(K_X + nD))$  cannot be greater than  $cn$  for  $c > 0$ . So

$$\frac{1}{12}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) = 0.$$

By Riemann-Roch,  $h^2(X, \mathcal{O}_X(nD)) = h^2(X, \mathcal{O}_X) - h^3(X, \mathcal{O}_X)$  is a constant and  $h^2(\mathcal{O}_D(nD)) = h^1(\mathcal{O}_D(nD))$  for all  $n \geq 0$ . Since  $h^2(X, \mathcal{O}_X) - h^3(X, \mathcal{O}_X) \geq 0$  and  $q = 0$ ,  $\chi(\mathcal{O}_X) \geq 1$ . By a theorem of Miyaoka [Mi],  $K_X$  is not nef.

Q.E.D.

**Proposition 3.6.** *Under the condition of Proposition 3.1, if  $\mathcal{O}_D(D)$  is torsion, then  $h^1(X, \mathcal{O}_X(nD))$  are bounded for all  $n$  and  $\frac{1}{12}(c_1^2 + c_2) \cdot D = \chi(\mathcal{O}_D) \geq 0$ .*

*Proof.* Let  $p$  be the least natural number such that  $\mathcal{O}_D(pD) = \mathcal{O}_D$ . Let  $G$  be the divisor defined by  $D|_D$ , then  $\kappa(G, D) = 0$ . By Iitaka [I1], page 35,  $h^0(D, \mathcal{O}_D(nD)) = 0$  if  $p$  does not divide  $n$ . Lemma 2.1-2.5 in [Ku] still hold in our case, so  $h^1(X, \mathcal{O}_X(nD))$  are bounded for all  $n \geq 0$ . Since  $h^3(X, \mathcal{O}_X(nD)) = 0$  and  $D^3 = D^2 \cdot K_X = 0$ , by Riemann-Roch, we have

$$\chi(\mathcal{O}_X(nD)) = 1 - h^1(X, \mathcal{O}_X(nD)) + h^2(X, \mathcal{O}_X(nD)) = \frac{1}{12}(c_1^2 + c_2)nD + \frac{1}{24}c_1c_2.$$

So  $\frac{1}{12}(c_1^2 + c_2)D \geq 0$  since  $h^1(X, \mathcal{O}_X(nD))$  are bounded for all  $n \geq 0$ .

By the short exact sequence

$$0 \longrightarrow \mathcal{O}_X((np - 1)D) \longrightarrow \mathcal{O}_X(npD) \longrightarrow \mathcal{O}_D(npD) = \mathcal{O}_D \longrightarrow 0,$$

we have

$$\chi(\mathcal{O}_D) = \chi(\mathcal{O}_X(npD)) - \chi(\mathcal{O}_X((np - 1)D)) = \frac{1}{12}(c_1^2 + c_2)D \geq 0.$$

Q.E.D.

We have proved Theorem 1.3.

By the standard classification theory of surfaces, if  $\mathcal{O}_D(D)$  is not torsion,  $D$  might be one of the following surfaces: (1) ruled surfaces over an elliptic curve; (2) bi-elliptic surfaces; (3) Abelian surfaces; (4) Elliptic surfaces.

## REFERENCES

- [AK] Altman, A.; Kleiman, S., Introduction to Grothendieck Duality Theory, Lecture Notes in Mathematics, Springer-Verlag, 1970.
- [Ba] Bădescu, L., Algebraic Surfaces, Universitext, Springer-Verlag, 2001.
- [Br] Bredon, G.E., Sheaf Theory, Graduate Texts in Mathematics, Springer-Verlag, 1997.
- [C] Cutkosky, S. D., Zariski decomposition of divisors on algebraic varieties, Duke Math. J. 53(1986), no. 1, 149-156.
- [CS] Cutkosky, S. D.; Srinivas, V. On a problem of Zariski on dimensions of linear systems. Ann. of Math. (2) 137 (1993), no. 3, 531-559.
- [FrQ] Friedman, R.; Qin, Z., On Complex surfaces diffeomorphic to rational surfaces. Invent. Math. 120(1995), no.1, 81-117.
- [Ful] Fulton, W., Intersection theory. Second edition. A Series of Modern Surveys in Mathematics 2. Springer-Verlag, Berlin, 1998.
- [GH] Goodman, J., Hartshorne, R., Schemes with finite-dimensional cohomology groups, American Journal of Mathematics, v. 91, Issue 1, 258-266, 1969.
- [GrH] Griffiths, P. and Harris, J., Principals of Algebraic Geometry, John Wiley & Sons, Inc., 1994.
- [H1] Hartshorne, R., Algebraic Geometry, Springer-Verlag, 1997.
- [H2] Hartshorne, R., Ample Subvarieties of Algebraic Varieties, Lecture Notes in Mathematics, 156, Springer-Verlag, 1970.
- [H3] Hartshorne, R., Local Cohomology, Lecture Notes in Mathe., 41, Springer-Verlag, 1967.
- [I1] Iitaka, S., Birational Geometry for Open Varieties, Les Presses de l'Universite de Montreal, 1981.

- [I2] Itaka, S., Birational Geometry of Algebraic Varieties, ICM, 1983.
- [I3] Itaka, S., Birational geometry and logarithmic forms, Recent Progress of Algebraic Geometry in Japan, North-Holland Mathematics Studies 73, 1-27.
- [KM] Kollár, J., Mori, S., Birational Geometry of Algebraic Varieties, Cambridge University Press, 1998.
- [Ku] Kumar, N. Mohan, Affine-Like Surfaces, Journal of Algebraic Geometry, 2(1993), 689-703.
- [L] Lazarsfeld, R., Positivity in algebraic geometry. I, II. Springer-Verlag, Berlin, 2004.
- [LZ] Luo, Tie; Zhang, Qi, Holomorphic forms on threefolds, preprint, 2003.
- [Ma] Matsuki, Kenji, Introduction to the Mori program. Universitext. Springer-Verlag, New York, 2002.
- [Mi] Miyaoka, Yoichi, The Chern classes and Kodaira dimension of a minimal variety. Algebraic geometry, Sendai, 1985, 449-476, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 1987.
- [N] Neeman, A., Ueda theory: Theorems and problems, Mem. Amer. Math. Soc., No.415 (1989), 1-123.
- [NS] Norguet, F.; Siu, Y.T. Holomorphic convexity of spaces of analytic cycles. Bull. Soc. Math. France 105, 191-223(1977).
- [Sa1] Sakai, F.,  $D$ -dimensions of algebraic surfaces and numerically effective divisors. Compositio Math. 48 (1983), no. 1, 101-118.
- [Sa2] Sakai, F., Weil divisors on normal surfaces. Duke Math. J. 51 (1984), no. 4, 877-887.
- [Se] Serre, J. P., Quelques problèmes globaux relatifs aux variétés de Stein, Collected Papers, Vol.1, Springer-Verlag(1985), 259-270.
- [Sim] Simpson, C., The construction problem in Kahler geometry, to appear in the volume "Different Faces of Geometry", International Mathematics Series, vol.3.
- [Siu1] Siu, Y.-T., Analytic sheaf cohomology groups of dimension  $n$  of  $n$ -dimensional noncompact complex manifolds. Pacific J. Math. 28 1969 407-411.
- [Siu2] Siu, Y.-T., Analytic sheaf cohomology of dimension  $n$  of  $n$ -dimensional complex spaces. Trans. Amer. Math. Soc. 143, 77-94(1969).
- [Ued] Ueda, T., On the neighborhood of a compact complex curve with topologically trivial normal bundle, J. Math. Kyoto Univ. 22-4, 583-607, 1983.
- [Uen1] Ueno, K., Classification Theory of Algebraic Varieties and Compact Complex Spaces, Lecture Notes in Mathematics, v.439, 1975, Springer-Verlag.
- [Uen2] Ueno, K., Algebraic Geometry 2, Sheaves and Cohomology, Translations of Mathematical Monographs, 197. American Mathematical Society, Providence, RI, 2001.
- [Za] Zariski, O., The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface, Ann. of Math. (2), 76(1962), 560-616.
- [Zh1] Zhang, J., Threefolds with vanishing Hodge cohomology, Trans. Amer. Math. Soc. 357 (2005), no. 5, 1977-1994.
- [Zh2] Zhang, J., There exist nontrivial threefolds with vanishing Hodge cohomology, to appear in Michigan Mathematical Journal.
- [Zh3] Zhang, J., On the  $D$ -dimension of a certain type of threefolds, submitted.
- [Zha] Zhang, Q., Global holomorphic one-forms on projective manifolds with ample canonical bundles. J. Algebraic Geometry 6 (1997), 777-787.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA  
*E-mail address:* zhangj@math.missouri.edu